

The α -minimum convex polygon as a relevant tool for isotopic niche statistics

Pauline Fey, Yves Letourneur, Silvere Bonnabel

► To cite this version:

Pauline Fey, Yves Letourneur, Silvere Bonnabel. The α -minimum convex polygon as a relevant tool for isotopic niche statistics. Ecological Indicators, 2021, 130, 10.1016/j.ecolind.2021.108048 . hal-03460350

HAL Id: hal-03460350 https://unc.hal.science/hal-03460350

Submitted on 1 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Contents lists available at ScienceDirect







journal homepage: www.elsevier.com/locate/ecolind

The α -minimum convex polygon as a relevant tool for isotopic niche statistics



Pauline Fey^a, Yves Letourneur^a, Silvère Bonnabel^{b,*}

^a UMR ENTROPIE (UR-IRD-IFREMER-CNRS-UNC), Université de la Nouvelle-Calédonie, BP R4, 98851 Nouméa cedex, New Caledonia
^b Institut de Sciences exactes et appliquées (ISEA), Université de la Nouvelle-Calédonie, BP R4, 98851 Nouméa cedex, New Caledonia

ARTICLE INFO

Keywords: Minimum convex polygon Non-parametric statistics Niche overlap Stable isotope analysis Trophic niche

ABSTRACT

Ecological (isotopic) niche refers to a surface in a two-dimensional space, where the axes correspond to environmental variables that reflect values of stable isotopes incorporated in an animal's tissues. Carbon and nitrogen stable isotope ratios (δ^{13} C- δ^{15} N) notably provide precious information about trophic ecology, resource and habitat use, and population dynamics. Various metrics allow for isotopic niche size and overlap assessment. In this paper, we advocate α -minimum convex polygons (MCP) - that have long been used for home range estimation – as a relevant tool for isotopic niche size, overlap, and characteristics. The method allows for outlier rejection while being suited to data that are not Gaussian in the bivariate isotopic (δ^{13} C- δ^{15} N) space. The proposed indicators are compared to other existing approaches and are shown to be complementary. Notably an indicator of divergence within the niche is introduced, and allows for comparisons at low (n > 6) and different sample sizes. The R code is made publicly available and will enable ecologists to perform isotopic niche comparison, contraction and expansion assessment, and overlap, based on various methods.

1. Introduction

Stable isotope analysis provides deep insight into a great variety of trophic and ecological processes. Since diet studies do not allow for a comprehensive view of food webs, it is necessary to use complementary methods. Isotope signatures in organisms yield information on time-integrated assimilated food (Fry, 1988; Vander Zanden and Rasmussen, 1999). Notably, carbon and nitrogen stable isotope ratios in the δ -space have been successfully used to understand the trophic dynamics in marine systems, to trace the pathway of organic matter of different origins through aquatic food webs (Fry and Scherr, 1984; Kaehler et al., 2000; Pinnegar and Polunin, 2000; Briand et al. 2016), to measure the impact of invasive species (O'Farrell et al., 2014), the impact of predators on preys (Gallagher et al., 2017), or to support the resource breadth hypothesis (Rader et al., 2017), to cite a few applications.

Regarding trophic food webs, carbon isotope composition in living animals usually provides indication of the origin of the ingested organic matter through a low increase in δ^{13} C per trophic level of about 1–1.4 ‰ on average (DeNiro and Epstein, 1978; Wada et al., 1991; Sweeting et al., 2007a). Nitrogen isotope ratio can be used as a proxy of the trophic level of organisms, as δ^{15} N usually increases to about 3.0–3.4 ‰

from food to consumer (Minagawa and Wada, 1984, Sweeting et al., 2007b). Thus, combined measurements of both isotopes can provide information on source material and trophic level, allowing for the construction of trophic relationships within the food web structure (Letourneur et al., 2013; Briand et al., 2016). Within a species, a genus, or a family, the space occupied by the individuals in the δ^{13} C *versus* δ^{15} N biplot is called isotopic niche, and can be used as an indicator of the trophic diversity of the species (or genus or family) (Newsome et al., 2007).

Layman et al. (2007) pioneered metrics to quantify isotopic niche structure. Ecologists routinely use their total area metric (TA), which is the area of the minimal convex polygon (MCP), or convex hull, containing all the organisms. Indeed, TA provides a useful ecological indication of the actual isotopic space occupied by a species or a community. In Jackson et al. (2011), the authors proposed Gaussian ellipses as an alternative to convex hulls, and introduced the indicator SEA which is the area of the theoretic confidence ellipse containing 40 % of a bivariate Gaussian having the same covariance matrix as the data. Both indicators were empirically compared (Syväranta et al., 2013). An extension of the ellipse method to *n*-dimensional niches has been proposed in Swanson et al., 2015, and recently an alternative method based

https://doi.org/10.1016/j.ecolind.2021.108048

Received 24 February 2021; Received in revised form 20 June 2021; Accepted 27 July 2021 Available online 31 July 2021

1470-160X/© 2021 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

^{*} Corresponding author. *E-mail address:* silvere.bonnabel@unc.nc (S. Bonnabel).



Fig. 1. Illustration of the 100 % MCP (light grey), and the 66% MCP (dark grey). Data randomly generated with Gaussian distribution (the right plot is obtained by contracting and dilating the left plot). The index TA₁ is the area of the total convex hull, whereas TA_{0.66} is the minimum convex polygon containing the most central 66% points, i.e., 33 out of 50 points.

on a kernel probability density function estimation of the niche has been introduced in Eckrich et al. (2020). Cucherousset and Villéger (2015) have also provided complementary isotopic niche indicators (isotopic divergence, dispersion, evenness and uniqueness). These indices are multidimensional, abundance-weighted and mathematically independent of the number of organisms analyzed. Niche metrics allow for comparisons between populations, see e.g., Andrades et al. (2019), notably through the study of overlap between occupied niches, see Botta et al. (2018), or to compare population isotopic niches between different habitats (Letourneur et al., 2017).

In behavioral ecology, for home range estimation (i.e., the habitat extent), the minimum convex polygon (MCP) is one of the most widely used methods. It may be traced back to Mohr (1947), and ecologists routinely use the 95% MCP instead, to rule out occasional sallies, see e.g, Van Beest et al. (2010). Inspiring from these works, we advocate the use of MCPs for the study of isotopic niches, as a basis for complementary and relevant indicators we introduce herein. We insist though, that the proposed indicators are devoted to isotopic niche assessment, not home range estimation. The interest will be shown hereafter using both simulated and experimental data in the field of marine biology.

The functions in the statistical environment R are made publicly available at http://www.silvere-bonnabel.com/TA_alpha. They are simple to use even for researchers who are not familiar with R, and allow for reproduction of the indicators and figures of the paper.

2. The TA_{α} index as a relevant tool

2.1. Definition of the TA_{α} index

Considering N points in the bivariate isotopic space, N being possibly small, the minimum convex polygon (MCP) of a given set of points is obtained by connecting the points that encompass the data. In other words, MCP is their convex hull. Moreover, it is also possible to define the α -MCP as the polygon of smallest area that encompasses a ratio α (hence between 0 and 100%) of the data. In turn, we define the index TA_{α} as the area of the α -MCP. TA_{α} is thus a value that represents an area, not to be confused with the α -MCP. Suppose $\alpha = M/N$. This means the α -MCP contains exactly M points by having eliminated the most exterior points. The proposed corresponding index TA_{α} is computed as follows.

• Find the convex hull of the N points. Amid the points that support the MCP, remove the one being the farthest to the barycenter of the polygon and recompute the MCP enclosing the remaining N-1 points.

- Repeat until the polygon contains M points.
- $\bullet\,$ Compute its area: it is equal to $TA_{M/N}.$

The user may set any value for α , thus not necessarily of the form M/N, especially when comparing samples with different sizes N. In the case where α falls strictly between two such ratios, i.e., $(M-1)/N < \alpha < M/N$ for some integer M, it is not obvious if the area computation, namely TA_{α}, should be based on the MCP containing M points or on the MCP containing M–1 points. Moreover, we intuitively want TA_{α} to continuously vary with α to avoid possible jumps in the index. To achieve this, we linearly interpolate between the area of both polygons, in function of the position of α in the interval [(M–1)/N, M/N] as follows.

• For $(M-1)/N < \alpha < M/N$, repeat above procedure until polygon contains M-1 points, and set TA_{α} = $(M-N\alpha) TA_{M/N} + (N\alpha-M + 1) TA_{(M-1)/N}$.

Fig. 1 illustrates the 100% and 2/3 (i.e. 66%) MCPs for two bivariate ($\delta^{13}\text{C-}\delta^{15}\text{N}$) plots.

2.2. Desirable features of the Ta_{α} index

The TA_{α} index inherits from the properties of the α -MCP, and generalizes the well-established TA indicator. In this respect, it possesses key properties that are as follows. First, the α -MCP contains the actual proportion α of data (up to a possible small discrepancy at low sample size we accommodate using interpolation) regardless of the underlying distribution of the data (no assumption is made that the distribution is Gaussian). Second, the indicator is robust to outliers as soon as α is not near 1, a feature that actually prompted the introduction of α -MCP in behavioral ecology, e.g., (Van Beest et al., 2010). Third, the indicator matches the boundedness of the support of actual ecological data, and coincides with the well-known TA indicator (Layman et al., 2007) in the extreme case $\alpha = 1$. This confirms its ecological relevance, in the sense that it reflects the size of an actually occupied isotopic niche space.

2.3. The TA_{α}/TA index

The TA index yields no indication of the distribution of the points within the convex hull. As complementary indicators, we introduce a family of indexes TA_{α}/TA that yields an indication of the variability within the convex hull, that is, divergence (see Cucherousset and Villéger, 2015). The TA_{α}/TA index is close to 1 if many points lie close to



Fig. 2. The effect of sample size N on the ratios SEA/TA, TA_{α}/TA and $(TA_{\alpha}/TA)^{c}$ for underlying bivariate normal centered distribution, with $\alpha = 2/3$, although other values are possible and lead to similar results. Results are averaged over 1000 Monte-Carlo runs.

the border of the convex hull, whereas it is low if most points are concentrated within a core area. The value of α may be arbitrarily set by the user. The value 2/3 is illustrative because it is based on simple proportions, but any other value is possible, as long as it is not too close to 0 or 1, in which case it may become meaningless. For example, if we find that TA_{2/3}/TA = 0.3 it means that a core group that contains two third of the population occupies a region as small as 30% of the total area. This may prove useful to study for instance the effect of the introduction of non-native species with extreme trophic positions (Cucherousset et al., 2012).

The proposed TA_{α}/TA possesses two desirable features. First, it is mathematically independent from the convex hull area, a feature it shares with the isotopic divergence ratio of Cucherousset and Villéger (2015), and not with previous metrics such as the mean nearest neighbor distance (Layman et al., 2007). This property is illustrated on Fig. 1, where the two plots actually display the same points, up to a (different) scaling factor along each axis. As scalings affect areas in the same proportions, the TA_{α}/TA ratio proves identical in both plots. In this sense, the indicator is independent of the total occupied area and only depends on the dispersion within the convex hull. Second, the proposed indicator is weakly sensitive to sample size N. This allows for comparison between samples having different sizes. This point is further developed in the next subsection.

2.4. The corrected $(TA_{\alpha}/TA)^{c}$ index

Albeit weakly sensitive to sample size, the ratio TA_{α}/TA drops at very low size, typically N \leq 10. To allow for isotopic niche comparison, we may inspire from the SEA_c methodology (Jackson et al., 2011), and correct the ratio at low sample size. To do so, we use a Monte-Carlo simulation to easily estimate the average value $\beta = \mathbb{E}\left(\frac{TA}{TA_{\alpha}}\right)$ of the ratio for a bivariate Gaussian distribution with sample size equal to N, and correct the TA_{α}/TA using β to make it unbiased with respect to sample size at low N. The corrected ratio $(TA_{\alpha}/TA)^c$ is displayed on Fig. 2. By contrast, SEA/TA represents the area of the 66% ellipse



	Cloud 1	Cloud 2		
TA _α	9.01	7.70		
Area intersect	5.03			
Area remaining	0.70			
TA_{α} both populations	12.37			

Fig. 3. Illustration of overlap between α -MCPs. The 95% ratio α allows for outlier rejection (4 points in this example). The R function allows for computation of the total convex hull encompassing both populations, as well as the area of the intersection and the area that remains outside the intersection. Data randomly simulated.



Fig. 4. Comparison between the α -MCP and the α – confidence ellipses for α taking the values (0.40, 0.66 and 0.95), for *Pomacentrus adelus* (N = 26) from New Caledonia (Briand et al., 2016) and *P. coelestis* (N = 40) from Marquesas Islands (Fey et al. in prep). The large dot in the center is the barycenter of the ellipses. The 40% ellipse captures only 33% of the population on the left plot, and as large as 50% on the right plot, *versus* 40% and 42%, respectively, for the TA_{rr}.

divided by TA, as used in Letourneur et al. (2017). Both indicators converged to the same value asymptotically, but SEA/TA is biased at sample sizes that may be of interest and thus may not serve to compare populations having different sizes.

2.5. The α -MCP overlap

When considering two groups, it may be interesting to compute both the actual common region of the space occupied by a ratio α of the members, that is, the area of the intersection between the two α -MCPs, as well as the convex hull of both α -MCPs. The former provides an indication of overlap whereas the latter represents a potential isotopic space. For instance, in some ecological contexts, overlap area might represent either a potential competition process or sharing for food resources between groups, whereas the whole α -MCP area might represent the entire isotopic space potentially available for the group of interest (Fig. 3). Using a value of α being less than 1 allows for outlier rejection, even for $\alpha = 0.95$, see Fig. 3.

3. Comparisons and discussion

In this section, we first briefly review the state-of-the-art ellipse methods (Jackson et al., 2011) as well as the recently introduced Kernelbased approach Eckrich et al., 2020. In a second step, we illustrate in the fourth subsection hereafter the relevance of the tools we have introduced using ecological data, collected on Pacific coral reef fish, and how they compare to those indicators and complement them.

3.1. The standard ellipse

The standard ellipse consists in assuming the distribution of the points to be Gaussian, then computing the empirical covariance matrix Σ , and applying a correction factor for unbiasedness with respect to sample size for bivariate Gaussians, namely $\Sigma \leftarrow \frac{N}{N-2}\Sigma$. Finally, the corresponding 40 % confidence ellipse, and its area, are computed. This yields the SEA_c indicator, and by omitting the correction factor N/(N-2) one recovers the SEA indicator.

Ellipse methods rely on an underlying Gaussian assumption whereas indicators based on MCPs are non-parametric. Moreover, the area of the confidence ellipse that theoretically contains a proportion α of the distribution is a pure dilatation/contraction of the 40% ellipse that is independent of the actual data. From an ecological viewpoint it may be

unsatisfying, especially for non-Gaussian underlying distributions, and it leads to erroneous values for large α , since the size of the ellipse tends to infinity when α approaches 1.

The SEA method was shown to suffer from uncertainty at sample size with less than 30 individuals, and caution is advised when populations are skewed (Syväranta et al., 2013).

3.2. Kernel-based methods

The idea of kernel-based estimation is to use a kernel to transform a sample into a continuous probability density function of a distribution. In turn, this allows for the calculation of the area encompassed by the contours a proportion α of data, e.g., 95%. The interest of Kernel-based methods is to more closely reflect the actual area occupied by the data, by relaxing the convex assumption for the contour, see Eckrich et al., 2020, hence offering versatility with respect underlying distributions as particularly desirable for home range study, (Getz et al., 2007). However, kernel-based methods are not available at low sample size, as the rKIN package does not apply to samples containing less than 10 individuals.

3.3. Comparisons using ecological data

3.3.1. Empirical confidence sets in practice

Fig. 4 displays the 0.40, 0.66 and 0.95 MCPs (note that those values were arbitrarily set and it is possible to choose other values) and confidence ellipses for two populations of coral reef fishes (Pomacentrus adelus and P. coelestis) sampled in SW lagoon of New Caledonia (Briand et al., 2016) and Marquesas' Islands, French Polynesia, respectively (unpublished data from Fey and Letourneur, in press). It evidences how the underlying true statistical distribution may differ from the bivariate Gaussian, as both look skewed. This is important as the ellipse of (Jackson et al., 2011) strongly relies on a Gaussian assumption. In both cases, the 95 % ellipses clearly appear too large. By contrast the α-MCPs inherently captures the correct amount of data points, up to a possible discrepancy of one individual as the value α does not fall exactly on a ratio M/N. The plots show a large gap between the theoretical and the actual proportion of the ellipse method. This advocates the use of α -MCPs as confidence sets that correctly capture the desired amount of data.

Table 1

Comparison of indicators for four coral reef fish species over healthy and degraded reefs in the SW New Caledonian lagoon (see Letourneur et al., 2017). N represents the number of individuals. TA_{α}/TA as an indicator of the shape of the distribution reveals larger discrepancies between both types of habitats than the other indicators. Moreover, its independence with respect to sample size makes it suited for comparisons.

		Chaetodon lunulatus		Chrysiptera rollandi		Halichoeres melanurus		Zebrasoma velifer	
		Degraded	Healthy	Degraded	Healthy	Degraded	Healthy	Degraded	Healthy
N		12	12	32	26	20	21	7	8
TA		0.56	1.01	0.99	0.88	1.40	1.29	0.21	0.57
SEAc/TA		0.46	0.47	0.34	0.36	0.45	0.35	0.96	0.70
Kernel/TA	0.40	0.43	0.43	0.43	0.45	0.65	0.36	-	-
	0.66	1.01	1.05	0.88	0.84	1.22	0.76	-	-
ΤΑ _α /ΤΑ	0.40	0.04	0.12	0.20	0.21	0.27	0.13	0.04	0.04
	0.66	0.25	0.39	0.49	0.46	0.61	0.38	0.10	0.28
(TA _α /TA) ^c	0.40	0.05	0.16	0.20	0.21	0.29	0.14	0.09	0.07
	0.66	0.26	0.40	0.49	0.45	0.61	0.37	0.11	0.30



Fig. 5. Kernel (left), TA_{α} (right). *G. chilospilus* (n = 133, red), *P. adelus* (n = 26, blue), *C. bathyphilus* (n = 7, green), at different α values (0.40, 0.66 and 0.95). Note that the Kernel based-approach cannot take into account the latter species case because n < 10.

3.3.2. The TA_a /TA as a way to measure discrepancies

Owing to the ability of TA_{α} /TA to compare populations with different sizes and areas (Fig. 2), this novel indicator is used as follows. In a recent article, Letourneur et al. (2017) investigated niche isotopic sizes of four coral reef fish species living in two different (degraded versus healthy coral reef) habitats from New Caledonia (see their Methods section for more details on sample collections, etc.). The indicator SEA_c/TA is therein used as a way to measure discrepancies between communities that lived in both coral reefs. We see that amongst all indicators, TA_{α} /TA, or its corrected version, indicates the most significant discrepancies between degraded and healthy reefs (Table 1) and as such better discriminates the statistical distributions. For instance, the variability for the Chaetodon lunulatus is identical in healthy and degraded reefs according to the SEA_c/TA indicator whereas it is by 36% lower using proposed TA_{α}/TA or its corrected version, with $\alpha = 0.66$ (Table 1). This might indicate the TA_{α}/TA is more accurate, as the result is more logical given the ecological requirements of that species, an obligatory corallivore (feeding on live coral). It is thus not surprising here, and even expected, to obtain a lower TA_a /TA ratio (or its corrected version) on the degraded reef, as the space occupied by most of the individuals tends to shrink. However, this expected result is not achieved when using a hybrid ratio instead that builds on a Kernel-based α -isopleth, see Table 1. This is logical, as the Kernel isopleth is less sensitive to sample size than TA (and TA_{α}), making the ratio Kernel/TA

more sensitive than TA_{α} /TA (see Fig. 2).

Results for *Zebrasoma velifer* are somewhat counterintuitive. Indeed, the area occupied shrinks in the degraded reef, although the species can be favored by reef degradation, as an herbivore. The fact that SEA_c/TA be much higher than TA_{0.40}/TA (Table 1), though, is in part an artefact due to SEA_c being corrected at low sample size, and TA being uncorrected. Although the corrected TA_α/TA index reflects a more concentrated niche in the degraded reef, as the space occupied by the most central members is in proportion lower, its unbiasedness at low sample size makes it more reliable, indicating that more complex ecological processes might be involved, or that the low size collected sample may merely not be representative.

3.3.3. Overlap indicators

Let us compare the proposed overlapping indicators based on α -MCPs with the Kernel-based approach (Eckrich et al., 2020). Three coral reef fish species having different sample sizes, collected in the SW lagoon of New Caledonia (*Gymnothorax chilospilus, Cirrhilabrus bathyphilus, Pomacentrus adelus*) (Briand et al., 2016) are displayed on Fig. 5. We see the interest of MCP is to leave some potential outliers out of the estimated niche (see two rightmost points), a feature not necessarily achieved with kernel-based methods. Indeed, the latter do not enforce convexity, allowing them to accommodate a large range of statistical distributions. However, in the context of marine ecosystems we believe

convexity makes sense as all values between the observed extremes are most likely biologically relevant, each point in the biplot being the result of a monthslong time-integrated diet where isotope quantities continuously evolve. It is thus not relevant to exclude values that lie in the biplot between two members, making the convex hull a domain that should be part of the area computation. For instance, considering the case of *C. bathyphilus*, isotopic values ranged from 6.64 to 7.71 ‰ for δ^{15} N and from -18.06 to -16.48 ‰ for δ^{13} C (Fig. 5, green dots), and it seems highly plausible that other unsampled individuals might possess C and/ or N isotopic values within these ranges (whereas there is no indication that values beyond this range are achievable by this group).

To conclude, two major points shall be noted. First, it was evidenced the proposed TA_{\alpha}/TA index, or its corrected version, does not suffer from some caveats of the other indicators, notably at low sample size. Moreover it may be used across the entire range of \alpha's, by contrast to the imposed choice of \alpha = 0.40 in the case of classical ellipses (Layman et al., 2007). Second, from an "ecological" point of view, it was evidenced through several examples of ecological interest that the proposed TA_\alpha/TA index, albeit simple and easily understandable to all, allowed for more interpretable results relative to the known biological requirements of the studied species.

CRediT authorship contribution statement

Pauline Fey: Data curation, Conceptualization, Methodology, Software, Writing - review & editing. **Yves Letourneur:** Conceptualization, Writing - review & editing, Supervision. **Silvère Bonnabel:** Conceptualization, Methodology, Software, Writing - original draft.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Andrades, R., Jackson, A.L., Macieira, R.M., Reis-Filho, J.A., Bernardino, A.F., Joyeux, J. C., Giarrizzo, T., 2019. Niche-related processes in island intertidal communities inferred from stable isotopes data. Ecol. Ind. 104, 648–658. https://doi.org/ 10.1016/j.ecolind.2019.05.039.
- Botta, S., Secchi, E.R., Rogers, T.L., Prado, J.H., de Lima, R.C., Carlini, P., Negrete, J., 2018. Isotopic niche overlap and partition among three Antarctic seals from the Western Antarctic Peninsula. Deep Sea Res. Part II: Topical Stud. Oceanogr. 149, 240–249. https://doi.org/10.1016/j.dsr2.2017.11.005.
- Briand, M.J., Bonnet, X., Guillou, G., Letourneur, Y., 2016. Complex food webs in highly diversified coral reefs: Insights from δ¹³C and δ¹⁵N stable isotopes. Food Webs 8, 12–22. https://doi.org/10.1016/j.fooweb.2016.07.002.
- Cucherousset, J., Blanchet, S., Olden, J.D., 2012. Non-native species promote trophic dispersion of food webs. Front. Ecol. Environ. 10 (8), 406–408. https://doi.org/ 10.1890/12.WB.018.
- Cucherousset, J., Villéger, S., 2015. Quantifying the multiple facets of isotopic diversity: new metrics for stable isotope ecology. Ecol. Ind. 56, 152–160. https://doi.org/ 10.1016/j.ecolind.2015.03.032.
- DeNiro, M.J., Epstein, S., 1978. Influence of diet on the distribution of carbon isotopes in animals. Geochimica et Cosmochimica Acta 42 (5), 495–506. https://doi.org/ 10.1016/0016-7037(78)90199-0.
- Eckrich, C.A., Albeke, S.E., Flaherty, E.A., Bowyer, R.T., Ben-David, M., Jackson, A., 2020. rKIN: Kernel-based method for estimating isotopic niche size and overlap. J. Anim. Ecol. 89 (3), 757–771. https://doi.org/10.1111/jane.v89.310.1111/1365-2656.13159.
- Fry, B., 1988. Food web structure on Georges Bank from stable C, N, and S isotopic compositions. Limnol. Oceanogr. 33 (5), 1182–1190. https://doi.org/10.4319/ lo.1988.33.5.1182.

- Fry, B., Scherr, E.B., 1984. 8¹³C measurements as indicators of carbon flow in marine and freshwater ecosystems. Contrib. Mar. Sci. 27, 13–47. https://doi.org/10.1007/978-1-4612-3498-2 12.
- Gallagher, A.J., Shiffman, D.S., Byrnes, E.E., Hammerschlag-Peyer, C.M., Hammerschlag, N., 2017. Patterns of resource use and isotopic niche overlap among three species of sharks occurring within a protected subtropical estuary. Aquat. Ecol. 51 (3), 435–448. https://doi.org/10.1007/s10452-017-9627-2.
- Getz, W.M., Fortmann-Roe, S., Cross, P.C., Lyons, A.J., Ryan, S.J., Wilmers, C.C., Coulson, T., 2007. LoCoH: nonparameteric kernel methods for constructing home ranges and utilization distributions. PLoS ONE 2 (2), e207. https://doi.org/10.1371/ journal.pone.0000207.
- Jackson, A.L., Inger, R., Parnell, A.C., Bearhop, S., 2011. Comparing isotopic niche widths among and within communities: Siber-stable isotope Bayesian ellipses in r. J. Anim. Ecol. 80 (3), 595–602. https://doi.org/10.1111/j.1365-2656.2011.01806.
- Kaehler, S., Pakhomov, E.A., McQuaid, C.D., 2000. Trophic structure of marine food web at the Prince Edward islands (Southern Ocean) determined by $\delta^{13}C$ and $\delta^{15}N$ analysis. Mar. Ecol. Prog. Ser. 208, 13–20. https://doi.org/10.3354/meps208013.
- Layman, C.A., Arrington, D.A., Montana, C.G., Post, D.M., 2007. Can stable isotope ratios provide for community-wide measures of trophic structure? Ecology 88 (1), 42–48. https://doi.org/10.1890/0012-9658(2007)88[42:CSIRPF]2.0.CO;2.
- Letourneur, Y., Lison De Loma, T., Richard, P., Harmelin-Vivien, M.L., Cresson, P., Banaru, D., Fontaine, M.-F., Gref, T., Planes, S., 2013. Identifying carbon sources and trophic position of coral reef fishes using diet and stable isotope (8¹⁵Nn and 8¹³C) analyses in two contrasted bays in Moorea, French Polynesia. Coral Reefs 32 (4), 1091–1102. https://doi.org/10.1007/s00338-013-1073-6.
- Letourneur, Y., Briand, M.J., Graham, N.A.J., 2017. Coral reef degradation alters the isotopic niche of reef fishes. Mar. Biol. 164 (12), 224. https://doi.org/10.1007/ s00227-017-3272-0.
- Minagawa, M., Wada, E., 1984. Stepwise enrichment of ¹⁵N along food chains: further evidence and the relation between 8¹⁵N and animal age. Geochimica et Cosmochimica Acta 48 (5), 1135–1140. https://doi.org/10.1016/0016-7037(84) 90204-7.
- Mohr, C.O., 1947. Table of equivalent populations of North American small mammals. Am. Midland Nat. 37 (1), 223–249. https://doi.org/10.2307/2421652.
- Newsome, S.D., Martinez del Rio, C., Bearhop, S., Phillips, D.L., 2007. A niche for isotopic ecology. Front. Ecol. Environ. 5 (8), 429–436. https://doi.org/10.1890/ 060150.1.
- O'Farrell, S., Bearhop, S., McGill, R.A.R., Dahlgren, C.P., Brumbaugh, D.R., Mumby, P.J., 2014. Habitat and body size effects on the isotopic niche space of invasive lionfish and endangered Nassau grouper. Ecosphere 5 (10), 1–11. https://doi.org/10.1890/ ES14-00126.1.
- Pinnegar, J.K., Polunin, N.V.C., 2000. Contributions of stable-isotope data to elucidating food webs of Mediterranean rocky littoral fishes. Oecologia 122 (3), 399–409. https://doi.org/10.1007/s004420050046.
- Rader, J.A., Newsome, S.D., Sabat, P., Chesser, R.T., Dillon, M.E., Martínez del Rio, C., Jackson, A., 2017. Isotopic niches support the resource breadth hypothesis. J. Anim. Ecol. 86 (2), 405–413. https://doi.org/10.1111/jane.2017.86.issue-210.1111/1365-2656.12629.
- Swanson, H.K., Lysy, M., Power, M., Stasko, A.D., Johnson, J.D., Reist, J.D., 2015. A new probabilistic method for quantifying n-dimensional ecological niches and niche overlap. Ecology 96 (2), 318–324. https://doi.org/10.1890/14-0235.110.1890/14-0235.1.sm.
- Sweeting, C.J., Barry, J.T., Polunin, N.V.C., Jennings, S., 2007a. Effects of body size and environment on diet-tissue δ¹³C fractionation in fishes. J. Exp. Mar. Biol. Ecol. 352 (1), 165–176. https://doi.org/10.1016/j.jembe.2007.07.007.
- Sweeting, C.J., Barry, J., Barnes, C., Polunin, N.V.C., Jennings, S., 2007b. Effects of body size and environment on diet-tissue δ¹⁵N fractionation in fishes. J. Exp. Mar. Biol. Ecol. 340 (1), 1–10. https://doi.org/10.1016/j.jembe.2006.07.023.
- Syväranta, J., Lensu, A., Marjomäki, T.J., Oksanen, S., Jones, R.I., 2013. An empirical evaluation of the utility of convex hull and standard ellipse areas for assessing population niche widths from stable isotope data. PloS One 8 (2), e56094. https:// doi.org/10.1371/journal.pone.0056094.
- Van Beest, F.M., Mysterud, A., Loe, L.E., Milner, J.M., 2010. Forage quantity, quality and depletion as scale-dependent mechanisms driving habitat selection of a large browsing herbivore. J. Anim. Ecol. 79 (4), 910–922. https://doi.org/10.1111/ j.1365-2656.2010.01701.x.
- Vander Zanden, M.J., Rasmussen, J.B., 1999. Primary consumer δ¹³C and δ¹⁵N and the trophic position of aquatic consumers. Ecology 80 (4), 1395–1404. https://doi.org/ 10.1890/0012-9658(1999)080[1395:PCCANA]2.0.CO;2.
- Wada, E., Mizutani, H., Minagawa, M., 1991. The use of stable isotopes for food web analysis. Crit. Rev. Food Sci. Nutr. 30 (4), 361–371. https://doi.org/10.1080/ 10408399109527547.